

EXHIBIT 6

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

CONSUMPTION OF OMEGA-3 FATTY ACIDS MAY REDUCE THE RISK OF CORONARY HEART DISEASE – A REVIEW

Mary G. Enig, Ph.D., F.A.C.N.
Nutritional Sciences Division
Enig Associates, Inc.
12501 Prosperity Drive, Suite 340
Silver Spring, Maryland 20904-1689 U.S.A.
Tel: 301-680-8600 Fax: 301-680-8100
15 November 1999

INTRODUCTION

Cardiovascular disease is reported as the leading cause of death in the United States. About 30 million Americans suffer from heart or blood vessel disease and as many as 27 million from high blood pressure. In 1991, 765,000 people died of heart disease and 190,000 died of stroke and other vascular complications--a total of 400 deaths per 100,000 people--and the number of deaths has increased since then. Cardiovascular disease accounts for 45 percent of all deaths in the United States. A quarter of the people who have a heart attack die within three hours of the first symptoms. Many experience sudden unexpected death and do not even reach a hospital. Another 25 percent die within the first few weeks after a heart attack. Cardiovascular disease is reported to cost the United States \$65 billion annually; more than \$38 billion of this cost is related to premature death (i.e., death before the age of 65).

Some dietary factors play a major role in determining the blood level of cholesterol, various lipoproteins, and triglycerides, which are considered major risk factors for cardiovascular disease. Some foods and individual nutrients appear to have a protective role in cardiovascular disease as well as in cancer.

The fatty acids eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) are omega-3 fatty acids found in most cold-water fish, such as tuna, salmon and mackerel, fish oils such as unhydrogenated menhaden oil, and fish liver oils such as cod liver oil. The adequate consumption of this class of fatty acids normalizes the levels of blood lipids such as triglycerides, reduces the tendency for blood-clot formation, and stabilizes the heartbeat against potentially fatal arrhythmias, especially under conditions of inadequate coronary blood flow, such as during a heart attack. Fish oil rich in omega-3 fatty acids is recognized as having cardioprotective actions.

The fatty acid alpha-linolenic acid (ALA) is an omega-3 fatty acid found in a few commonly available unprocessed oils, especially flax seed oil, canola oil, soybean oil, and walnut oil and in other oils such as perilla oil. Refining such as deodorization and partial hydrogenation causes loss of the omega-3 fatty acid ALA, which can range from a minor loss to near total loss. When levels of this omega-3 fatty acid in the diet are not swamped by high levels

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

of the fatty acid linoleic acid (LA), an omega-6 fatty acid, i.e., the ratio of LA/ALA is not greater than 3.0, there is some conversion in the human from ALA to EPA and DHA, affording the body some accumulation of EPA and DHA in the phospholipids of circulating platelets and the protection it offers against platelet aggregation. Without adequate EPA and DHA in phospholipids, the modulating role of the derivative prostaglandins are not seen.

Omega-3 fatty acids are precursors to eicosanoids, e.g., prostaglandins and leukotrienes. These eicosanoids modulate metabolic actions in many tissues by their effects on, e.g., tissue hormones and prehormones, signal molecules, and receptors. They are responsible for maintenance of a desirable balance of, e.g., endogenous vasoconstriction, activity of the renin-angiotensin system, salt and water balance, peripheral sympathetic tone, cardiac output and baroreceptor settings, all functions in the body that are of critical importance to proper maintenance of a healthy cardiovascular system.

Adequate consumption of omega-3 fatty acids may reduce the risk of coronary heart disease through a number of effects on cardiovascular risk factors, especially related to reduction of triglycerides, improvement of heart rate variability, and risk of fatal arrhythmias. Support for this statement may be found in the following review and evaluation of recent scientific literature, which includes a discussion of:

- a) the competence and reliability of recent (1994-present) science supporting the claim;
- b) the level of intake required to achieve beneficial results and anticipated levels of omega-3 fatty acid consumption by use of a supplement compared to the probable level needed when the nutrient may be consumed in a well-balanced diet;
- c) the safety of the nutrient at dose levels and consumption levels needed to produce the physiologic effect.

EVALUATION OF CURRENT SCIENTIFIC LITERATURE WITH RESPECT TO THE EFFECT OF OMEGA-3 FATTY ACIDS ON CORONARY HEART DISEASE RISK FACTORS

In order to determine the extent to which the current scientific literature supports the claim that "consumption of omega-3 fatty acids may reduce the risk of coronary heart disease," 168 published articles were identified from Medline for the years 1994 to 1999 as relevant to the topic.

Of the 168 articles, 74 were reports of clinical treatments and clinical trials (44%), 16 were reports of epidemiological studies (9.5%), 36 were review/tutorial articles (21.4%), 28 were reports of basic research in animal and/or cell culture models (16.7%), 5 were letters to the editors (3%), 5 were articles in medical news letters (3%), and 4 were editorials (2.4%).

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

Of the reports identified by Medline as clinical treatments and clinical trials, 12 were published in 1999 (through September), 12 were published in 1998, 13 were published in 1997, 13 were published in 1996, and 12 were published in 1995. On examination several reports listed by Medline as clinical trials turned out to actually be review articles where the author included summary reports of clinical trials. In each case, the original publications from the trials were consulted for verification of the information.

CLINICAL TREATMENT AND CLINICAL TRIALS LITERATURE

Review of selected representative literature

Investigators of the **GISSI-Prevenzione trial (1999)** concluded that treatment with 1 gram per day of n-3 PUFA in the form of ethyl esters of eicosapentaenoic acid and docosahexaenoic acid (850-882 mg) "led to a clinically important and statistically significant benefit" of lowered the risk of the primary endpoint of a fatal cardiovascular event. This trial was reviewed for the Cardiovascular News Meeting Highlights of the 48th Scientific Sessions of the American College of Cardiology under the section on Preventive Cardiology by Ferguson (1999). The summary stated: "In patients with a recent history of MI, dietary supplementation with n-3 PUFA (1 g/day)...seems to significantly reduce long-term cardiac events, total mortality, and sudden cardiovascular death."

In this review, which Medline lists as a clinical trial publication, **Harris (1999)** gives an overview of the history of lipid-altering effects of the n-3 PUFA, an evaluation of low dose studies, a brief review of animal studies, and reports on two clinical intervention studies with n-3 PUFA and coronary heart disease, which were both prospective, randomized trials with hard endpoints, i.e., cardiac events, and mortality. The first trial, reported in 1989 (Harris' reference 8) was called DART, and used an intake of oily fish for the experimental group, which provided about 200 to 400 grams of fish a week and approximately 500 to 700 mg/day of eicosapentaenoic acid and docosahexaenoic acid. In the ensuing 2 years, those on the fish diet had 29% fewer deaths. The second trial is Singh et al (1997), known as the Indian Experiment. In this trial, approximately 360 patients who presented with myocardial infarction were randomized to 2 g of n-3 PUFA (6 capsules of Maxepa), 2.9 g of alpha-linolenic acid (20 g of mustard seed oil), or placebo. The patients were followed for 12 months with a significant reduction in new cardiac events. Harris concludes in summary that "there is extensive and consistent documentation of the ability of marine n-3 [PU]FA to reduce serum triglycerides levels in humans by 20-30% at doses of 3-4 g per day." Harris had previously reviewed several earlier studies of the impact of fish oil omega-3 on hypertriglyceridemia (see reviews section below, Harris 1996).

Based on an evaluation of 82 male children from age 7-14 years, who were divided into groups with and without risk factors of cardiovascular pathology **Kozycheva and Slezka (1998)** report a "correcting effect" by dietary omega-3 PUFA from marine sources on the composition of "erythrocyte fatty acid defects" at early stages of disease. This was especially observed in the case of children who had myocardial infarction heredity. The authors conclude that dietary

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

omega-3 PUFA can be used for preventative treatment of children with cardiovascular disease heredity.

Pogozheva et al (1998) conducted a clinical trial to study levels of natural antibodies against catecholamines in 138 patients with cardiovascular diseases. A comparative analysis of the influence of "antiatherosclerotic" diets with omega-3 PUFA from both plant and animal sources was made. The greatest "immunocorrection effect" was found when the diet was enriched with omega-3 PUFA freshwater fish fat.

Heart rate variability and omega-3 PUFA

Decreased heart rate variability (HRV) is associated with increased coronary mortality due to sudden cardiac death (Christensen JH et al 1999a), and in fact, HRV is considered "a powerful predictor of mortality and of arrhythmic events in humans." Findings from a number of clinical trials from several research facilities published between 1996 and 1999 reported on the beneficial effect between dietary intake of omega-3 polyunsaturated fatty acids and HRV. These research reports include: Christensen JH et al 1999a, Christensen JH et al 1999b, Christensen JH et al 1999c, Christensen JH et al 1998, Christensen JH et al 1997a, Christensen JH et al 1997b. A representative sample of these trials are reviewed.

Because there was evidence for an anti-arrhythmic effect of n-3 polyunsaturated fatty acids (n-3 PUFA) in animals, Danish researchers decided to evaluate a possible beneficial anti-arrhythmic effect of consumption of the long-chained n-3 PUFA on heart rate variability. Several studies were undertaken (Christensen JH et al 1999a). Three different groups of people were evaluated in these studies, and they included 1) patients with previous myocardial infarction (Christensen et al 1999a, 1997, 1996), 2) patients with chronic renal failure (Christensen JH et al 1998), and 3) healthy volunteers (Christensen JH et al 1999a). Additionally, Christensen JH et al (1999c) had validated the measures of HRV and risk for sudden cardiac death. The authors' conclusion was that "the results showed a beneficial effect of n-3 PUFA on HRV in...[these]...populations...[and that their]...data may explain the findings from other studies, namely that a low occurrence of sudden cardiac death is observed in subjects with and without coronary heart disease who regularly eat fish."

Leaf et al (1999) have reviewed the recent studies which they report identifies what is likely the major mechanism by which n-3 PUFA exerts its antiarrhythmic effect. They report that PUFA stabilize the electrical activity of isolated cardiac myocytes by inhibiting sarcolemmal ion channels; this then requires a stronger electrical stimulus to elicit an action potential resulting in a markedly prolonged refractory period.

An intervention trial was assessed in 52 patients with a history of previous myocardial infarction and left ventricular dysfunction. The content of n-3 PUFA in platelets was monitored and was found to be closely associated with the patient's fish-consuming habits. This study reported that a significant positive correlation was observed between the n-3 fatty acid docosahexaenoic acid and heart rate variability (Christensen et al 1997a).

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

The post-MI patients in the above study were randomly evaluated in the next study, which investigated the effect of supplementing dietary n-3 PUFA on ventricular arrhythmias and heart rate variability (HRV) in patients with a previous myocardial infarction. In a double-blind, placebo-controlled study, patients were given 5.2 g per day of n-3 PUFA or placebo and HRV response was measured from a 24 hour Holter recording both at baseline and, following supplementation, at the end of 12 weeks. The study showed significantly increased HRV after n-3 PUFA compared to both baseline values ($p = 0.04$) and to controls ($p = 0.01$). This was interpreted as supporting the hypothesis that n-3 PUFA may have an anti-arrhythmic effect in humans (Christensen et al 1997b)

In a clinical trial published in the *American Journal of Clinical Nutrition*, Christensen JH et al (1999a) studied the dietary intake of capsules containing 6.6 g or 2.0 g per day of long-chain n-3 polyunsaturated fatty acids (PUFA) reesterified from fish oil. Function was determined by a 24 hour Holter recording in healthy volunteers; this double-blind study included both males and females. The researchers' conclusion was that the "...study showed a beneficial effect of n-3 PUFA on HRV in healthy men, suggesting an antiarrhythmic effect of n-3 PUFA." The effect was not seen in the women.

Li et al (1999), evaluated thrombotic risk factors in vegetarian men who are known to have higher platelet aggregability than omnivores. The authors concluded that, in this study where the normal vegetarian diet was supplemented with added sources of ALA from vegetable oils such as canola and linseed, there was a significant increase in EPA and DHA as well as an increase in the ratio of n-3 to n-6 PUFAs in platelet phospholipids and plasma lipids in these subjects.

Singh et al (1997) (as reviewed below) compared the effects of both fish oils supplements and mustard oil in patients with suspected acute myocardial infarction. In this randomized, placebo-controlled trial, the effects of treatment with fish oil (eicosapentaenoic acid, 1.08 g/day) and mustard oil (alpha-linolenic acid, 2.9 g/day) were compared for 1 year. There were 122 patients treated with fish oil (group A), 120 patients treated with mustard oil (group B), and 118 patients treated with placebo (group C) all with suspected acute myocardial infarction (AMI). Treatments were started on average approximately 18 hours after the symptoms of AMI brought them in for treatment in all three groups. The extent of cardiac disease, rise in cardiac enzymes, and lipid peroxides were comparable among the groups at entry into the study. After 1 year total cardiac events were significantly less in the fish oil and mustard oil groups compared with the placebo group (24.5% and 28% vs. 34.7%, $p < 0.01$). Nonfatal infarctions were also significantly less in the fish oil and mustard oil groups compared with the placebo group (13.0% and 15.0% vs. 25.4%, $p < 0.05$). There was no significant reduction of total cardiac deaths in the mustard oil group but the fish oil group had significantly less cardiac deaths compared with the placebo group (11.4% vs. 22.0%, $p < 0.05$).

In a multicenter study, Archer et al (1998) have evaluated dietary intake of omega-3 fatty acids related to hemostatic factors which are known to play an important role in the complications of ischemic heart and vessel disease. Dietary fats such as omega-3 PUFA have

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

been shown to possibly influence hemostatic factors. It was noted that most studies reporting an inverse association between cardiovascular disease and fish and n-3 fatty acid consumption have used supplemental doses of fish oil or intakes exceeding the typical amount consumed by the U.S. population. This report examined the associations of usual intakes of fish, linolenic acid, eicosapentaenoic acid, and docosahexaenoic acid with fibrinogen, factor VII, factor VIII, and von Willebrand factor in the Coronary Artery Risk Development in Young Adults (CARDIA) Study. The analyses were reported for 1672 black and white men and women aged 24 to 42 years in 1992 to 1993. After adjustment for age, body mass index, diabetes, number of cigarettes smoked per day, race, and energy and alcohol consumption, no significant associations were observed between those who consumed no fish versus those who consumed the highest level of dietary fish with respect to fibrinogen, factor VIII, or von Willebrand factor for any race-sex group. Comparisons of tertile 1 versus tertile 3 for dietary linolenic acid, eicosapentaenoic acid, and docosahexaenoic acid were also not significantly associated with fibrinogen, factor VII, factor VIII, or von Willebrand factor for any race-sex group. These data were interpreted to suggest that customary intakes of fish and n-3 fatty acids in populations that generally do not consume large amounts of these food items are not associated with these hemostatic factors.

EPIDEMIOLOGICAL LITERATURE

Guallar et al (1999) studied the levels of omega-3 fatty acids in adipose tissue related to risk of myocardial infarction as part of the EURAMIC study. They found a slight protective effect of alpha-linolenic acid and no protective effect of docosahexaenoic acid and did not identify any eicosapentaenoic acid content.. Since, there are usually no elongated omega-3 fatty acids (omega-3 PUFA) stored in adipose, because they are basically membrane fatty acids, it is unlikely that the findings in this study have much meaning regarding the relationship between omega-3 PUFA and risk of myocardial infarction.

Kaminskas et al (1999) compared abdominal adipose fatty acids from two populations with different cardiovascular risk. Lithuanian men, who have four times higher coronary heart disease mortality than Swedish men, in spite of having lower total and LDL cholesterol, were found to have significantly higher percentage of long-chain PUFA and lower percentage of saturated fatty acids, especially myristic acid. Since the PUFA normally would be found in membrane tissue where they provide precursor to eicosanoids, and the saturates would normally be found in the adipose, it could be that the findings in this study may be indicative of some lipid abnormality unrelated to the long-chain PUFA specifically.

Hu et al (1999) evaluated the risk of fatal ischemic heart disease (IHD) among women as it was related to their dietary intake of alpha-linolenic acid. This prospective cohort study with ten years of follow-up was reported to support the hypothesis that a higher intake of alpha-linolenic acid is protective against fatal ischemic heart disease (IHD). Higher consumption of foods such as oil-based salad dressing that provide polyunsaturated fats, including alpha-linolenic acid, may play a role in the reduction of the risk of fatal IHD.

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

Aschereio et al (1996) examined the association between fat intake and the incidence of coronary heart disease in a cohort of men of middle age and older as part of the health professionals follow up study in the United States. The data did not support the strong association between intake of saturated fat and risk of coronary heart disease suggested by international comparisons, but it did support a specific preventive effect of linolenic acid intake.

REVIEWS

Roche and Gibney (1999) have reviewed studies which used both short- and long-term supplementation of omega-3 fatty acids in amounts ranging from 1 g per day to 28 g per day from fish oils. After evaluating the studies which showed lowering of both fasting and postprandial triglycerides, these researchers concluded that although "the precise biochemical nature of this effect remains to be established... reduction of postprandial plasma TG concentrations by n-3 PUFA may partly explain why n-3 PUFA intake is inversely related to CHD mortality.

Schmidt and Dyerberg (1999) have reviewed a number of trials and have concluded that a 1989 study provided "clinical evidence for a protective role of n-3 PUFA in CHD." They also note that the research (Christensen 1996) on heart rate variability showed a benefit from n-3 PUFA. However, in this review, the authors still call for increased trials for both fish and fish oil concentrates. They note that many populations do not have fish intake that would provide 1 g/week of n-3 PUFA, and that, additionally, "some people dislike seafood."

Connor and Connor (1997) ask the question "Are fish oils beneficial in the prevention and treatment of coronary artery disease?" Their answer is that the "n-3 fatty acids of fish and fish oil have great potential for the prevention and treatment of patients with coronary artery disease." They report that unlike many of the pharmaceutical agents used to treat coronary artery disease that have just a single mechanism of action, the omega-3 PUFA of fish oil have "multifaceted actions." As noted by other researchers, one of the most important effects of the omega-3 PUFA is the prevention of arrhythmias. The documentation for this effect has been derived from experiments in cultured myocytes, experiments in animals, epidemiologic correlations, and clinical trials. The Connors note that the ability of these omega-3 PUFA to inhibit ventricular fibrillation and consequent cardiac arrest is especially important. They discuss several mechanism including the several antithrombotic actions of EPA, particularly in inhibiting the synthesis of thromboxane A₂, the prostaglandin that causes platelet aggregation and vasoconstriction. Further they report that fish oil retards the growth of the atherosclerotic plaque by inhibiting both cellular growth factors and the migration of monocytes. Also, omega-3 PUFA is reported to promote the beneficial synthesis of nitric oxide in the endothelium. In experiments in humans, a profound hypolipidemic effect of fish oil, is seen, especially with regard to lowering of plasma triacylglycerol, and inhibition of very-low-density lipoprotein production and apolipoprotein B synthesis. Fish oil is also said to have a mild blood pressure-lowering effect in both normal and mildly hypertensive individuals. The Connors state in their abstract that "these composite effects suggest a prominent therapeutic role for fish oil in the prevention and treatment of coronary artery disease."

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

Harris (1996, 1999) has reviewed several studies of the impact of fish oil omega-3 on hypertriglyceridemia. He notes that triglyceride lowering is the most consistent effect of dietary fish oils, having been observed in Greenland Eskimos and subsequently in controlled clinical trials. In those early studies, total cholesterol levels are not altered, but there was a transient rise in LDL levels, especially in hypertriglyceridemic patients. In later studies which followed large numbers of patients from 6-9 months there was a persistence of the hypotriglyceridemic effect as long as the fish oil treatment continued, and there was evidence from the studies that the LDL raising effect may not persist. Further, Harris notes the safety of fish oils was also supported in the studies because problems with excessive bleeding and worsening glycemic control did not materialize. However, in one of the studies, dietary fish oil was reported to be ineffective in slowing the rate of restenosis after coronary angioplasty.

De Deckere et al (1998) reports on conclusions from an expert workshop which reviewed the health effects of n-3 polyunsaturated fatty acids (PUFA). Among their conclusions were that 1) consumption of fish may reduce the risk of coronary heart disease (CHD) and that people at risk for CHD are therefore advised to eat fish once a week. They noted that the n-3 PUFA in fish are probably the active agents in this effect. Therefore, people who do not eat fish should consider obtaining 200 mg of very long chain n-3 PUFA daily from other sources. They also felt there is incomplete but growing evidence that consumption of the plant n-3 PUFA, alpha-linolenic acid, reduces the risk of CHD and that an intake of 2 g/d or 1% of energy of alpha-linolenic acid appears prudent. They further noted that the ratio of total n-3 over n-6 PUFA (linoleic acid) is not useful for characterizing foods or diets because plant and marine n-3 PUFA show different effects, and because a decrease in n-6 PUFA intake does not produce the same effects as an increase in n-3 PUFA intake. Separate recommendations for alpha-linolenic acid, marine n-3 PUFA and linoleic acid were preferred.

ANIMAL AND CELL CULTURE STUDIES

An animal study by **Nair et al (1999)** found a response in the pig model that suggested a "possible role of nonesterified (n-3) polyunsaturated fatty acids in the prevention of arrhythmias."

Hishinuma T et al (1999) studied the effect of omega-3 PUFA on changes of lipid profiles and prostacyclin production by cultured bovine aortic endothelial cells. They report that their findings "suggest that EPA, docosapentaenoic acid, and DHA are interconverted to each other, and anti-aggregatory effects of EPA or DHA may be partially due to the stimulation of prostacyclin formation in endothelial cells."

Animal studies have confirmed the protective effect against ischemia-induced fatal arrhythmias (**Billman et al 1999**).

Hartman (1995) has reviewed the results of a 5-year study of survivors of a first myocardial infarction, which reports the prevention of secondary coronary events as an effect of

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

an alpha-linolenic acid-enriched diet (experimental) compared with the prudent diet of the American Heart Association (control). The data were from the dietary intervention study of de Lorgeril et al (1994). During an average of 27 months of follow-up, there were 5 nonfatal myocardial infarctions and 3 cardiac deaths in the experimental group versus 17 nonfatal myocardial infarctions and 16 cardiac deaths in the control group. The overall mortality was 20 in the control group and 8 in the experimental group. The interpretation of this study was that a diet enriched with alpha-linolenic acid appears to be effective in secondary prevention of coronary events, but the actual trial had three dietary components being used concurrently; in addition to the increased ALA component, the second component was increased consumption of fish, and the third component was that it was to be a "Mediterranean" diet.

Mantzioris et al 1995 reports that increased concentrations of cellular eicosapentaenoic acid (EPA) have been shown to be beneficial in coronary heart disease, hypertension, and inflammatory disorders. Successful long-term strategies for increasing cellular EPA concentrations require an understanding of the relationships between cellular concentrations of EPA and dietary amounts of ALA, a precursor of EPA, and dietary amounts of linoleic acid (LA), which can act as an antagonist of ALA conversion to EPA (and DHA). With this dietary intervention trial, which studied healthy human volunteers and which incorporated ALA-rich vegetable oil (flaxseed oil) against a background diet low in LA, the researchers were able to examine these relationships. Linear relationships were reported between dietary ALA and EPA in plasma fractions and in cellular phospholipids. By contrast there was no relationship observed between dietary LA and tissue concentrations of its metabolite, arachidonic acid (AA). There was an inverse relationship between dietary ALA and DHA concentrations in the phospholipids of plasma, neutrophils, mononuclear cells, and platelets. The results of this study indicated that increasing dietary ALA would elevate tissue EPA concentrations in a predictable manner. It was concluded that this insight could facilitate the rational planning of practical dietary strategies for the long-term elevation of EPA concentrations in tissues.

Santos et al 1995 studied the effects of dietary supplementation with fish on plasma fatty acid levels in 20 coronary heart disease patients who had suffered acute myocardial infarction. The study was divided into three periods which included the term of hospital admission, after 8 weeks on a special diet designed for patients with ischemic heart disease, and after 4 weeks on an n-3 PUFA-supplemented diet in which red meat was replaced with lean and fatty fish. At the end of each period, the subjects were given a 48-hour recall dietary questionnaire, to assess compliance with the diet, and blood samples were collected for the determination of plasma fatty acids. Stearic acid was significantly decreased after the fish diet, but fatty acids such as palmitoleic and oleic acid showed no significant changes throughout the study. At the end of the 4-week period when the fish diet was consumed there was a significant decrease in linolenic acid and the significant increase in its long-chain derivatives EPA and DHA in the plasma fatty acids, which was thought to show beneficial effects on coronary heart disease.

Horrocks and Yeo (1999) have reviewed the epidemiological studies which have shown a strong correlation between fish consumption and reduction in sudden death from myocardial infarction and the reduction is approximately 50% with 200 mg a day of DHA from fish.

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

DHA is considered the active component in fish. They note that not only does fish oil reduce triglycerides in the blood and decrease thrombosis, but it also prevents cardiac arrhythmias. They also suggest that the association of DHA deficiency with depression is the reason for the robust positive correlation between depression and myocardial infarction. Patients with cardiovascular disease are often advised to adopt a low-fat diet with a high proportion of carbohydrate, as are patients with Type II diabetes. However, a study with women showed that this type of diet increased plasma triglycerides and the severity of Type II diabetes and coronary heart disease. Both DHA and EPA are present in fatty fish (salmon, tuna, mackerel). DHA is also present at low levels in meat and eggs. The shorter chain n-3 fatty acid, ALA, noted not to be converted very well to DHA in man.

CONVERSION OF ALA TO EPA AND DHA IN HUMANS

There was previously thought to be substantial conversion of the omega-3 fatty acid alpha-linolenic acid (ALA) to the elongated omega-3 fatty acids eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), which are found most commonly in fish and fish oils. In fact, Bordoni et al (1996) had reported a 90% conversion of ALA to elongated forms of n-3 PUFA. Research reported in 1998 (Gerster), which measured the conversion of ALA labeled with radioisotopes, has shown that the conversion in adult humans is only approximately 6% to EPA and 3.8% to DHA when the background diet is high in saturated fat, but the conversion is reduced 40 to 50% when the diet is rich in omega-6 PUFA. Since the typical diet in the United States is high in omega-6 PUFA, the levels of EPA and DHA that can be obtained from ALA-rich vegetable oils through conversion in humans is not particularly high and are probably on the order of 3% conversion to EPA and less than 2% to DHA. In addition, most of the vegetable oils that have ALA only in their unprocessed form have had the biologically available ALA totally removed by partial hydrogenation or changed to a form that is not biologically available by refining such as deodorization and/or decolorization.

Suzuki et al (1997) measured the uptake into brain tissues of omega-3 fatty acids from different sources and provided an example of the extent to which ALA may be bioavailable. The percentage of DHA found in the brain was ranked in the following order: the salmon oil diet group > the sardine oil diet group > > the perilla oil diet group > > the lard and palm oil diet groups for the 30 day feeding trial. When the animals were fed purified fatty acids, the uptake was in the order of the DHA diet group > > the eicosapentaenoic acid and alpha-linolenic acid diet groups for the 6 day feeding trial.

IN SUMMARY:

THE KNOWN EFFECTS OF OMEGA-3 PUFA ON CORONARY HEART DISEASE

From a review of the current scientific literature one can see that some of the mechanisms have been identified whereby omega-3 PUFA from fish oil can be said to have beneficial physiological and biochemical effects, which result in the prevention of coronary heart disease. At least three such mechanisms, i.e. triglyceride lowering, the raising of heart rate variability,

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

and reduction of fatal arrhythmias have been well studied. In the studies reviewed, the adequate dietary intake of omega-3 PUFA, was primarily from whole fish oils.

In conclusion, it is reported in peer reviewed scientific literature that the dietary intake of adequate amounts of omega-3 PUFA has been shown to 1) decrease adversely high levels of triglycerides (TG) and very low density lipoproteins (VLDL), thus decreasing the risk of coronary heart disease; 2) increase heart rate variability and thereby decrease the risk of sudden death from heart attack thus helping to prevent a first heart attack; and 3) help prevent second heart attack in individuals who have sustained a first heart attack.

REGARDING LEVELS OF OMEGA-3 PUFA THOUGHT TO BE BENEFICIAL

The levels of omega-3 PUFA in the daily diet needed to maintain optimal omega-3 PUFA in the tissues and to provide physiologically and biochemically beneficial effects are approximately 3 to 4 g/day of fish oil omega-3, but some studies have reported benefits at 1 g/day omega-3 PUFA.

Levels of fish required to achieve the 3-4 g/day are much higher than the levels typically consumed in the United States. Not all fish have sufficiently high levels of fat to provide even 1 g/day in a typical 100 g serving.

Plant omega-3 (ALA) is variably converted to omega-3 PUFA depending on background diet of the individual; a background diet high in saturated fat is needed for optimal conversion and a diet high in omega-6 linoleic acid (LA) decreases conversion by about 50%. In the United States and Canada and other parts of the world this high saturated fat diet is discouraged and usually replaced with a diet high in omega-6 LA.

REPORTED LEVELS OF OMEGA-3 PUFA FROM FISH OILS AND OMEGA-3 ALPHA-LINOLENIC ACID FROM VEGETABLE OILS FED TO HUMANS

GISSI clinical trial (1999): 1g/d omega-3 PUFA long term containing 850-882 mg EPA and DHA as ethyl esters.

Roche and Gibney (1999) reviewed studies which used both short- and long-term supplementation ranging from 1 g/d omega-3 PUFA (Roche and Gibney 1996) to 28 g/d omega-3 PUFA (Harris et al 1988)

Harris (1999) reviewed "low-dose" 3-4 g/d omega-3 PUFA on average, and other studies "up to 25 g/d omega-3 PUFA

Tinker et al (1999) fed 13.2 g fish oil which contained 5.2 g omega-3 PUFA in a clinical study which reported significant effects on lowering postprandial triglyceride and lipoprotein levels.

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

von Schacky et al (1999) fed 3 g/day of fish oil which contained 1.5 g omega-3 PUFA for 21 months after an initial 3 months of 6 g/day of fish oil. They reported modest effects on coronary atherosclerosis with fish oil treated patients having fewer cardiovascular events.

Nordoy et al (1998) fed 4 g/day of purified omega-3 PUFA in conjunction with Simvastatin to patients with combined hyperlipidemia. They report that the combined treatment had a beneficial additive effect.

Sirtori et al (1997) fed (2.58 g x 3) 7.74 g/d of EPA and DHA as ethyl ester for 2 months and then (1.72 g x 2) 3.44 g/d of EPA and DHA as ethyl ester for an additional 4 months.

LEVELS OF OMEGA-3 PUFA IN FISH

According to **Connor and Connor (1997)** (Table 2 in *AJCN*), fish having less than 0.5 g omega-3 PUFA/100 g edible portion are carp, catfish, Atlantic cod, Pacific cod, flounder, haddock, halibut, ocean perch, walleye pike, red snapper, sole sturgeon, swordfish, brook trout, all crustaceans, New Zealand abalone, clam, octopus, scallop, squid. Fish having 0.5 to 1 g omega-3 PUFA/100 g edible portion are striped bass, Florida pompano, shark, rainbow trout, tuna, blue mussel, Pacific oyster. Fish having 1 to 1½ g omega-3 PUFA/100 g edible portion are European anchovy, carp, mullet, sablefish, all salmon. Fish having 1.5 to 2 g omega-3 PUFA/100 g edible portion are Atlantic herring, Pacific herring, lake trout. Fish having greater than 2 g omega-3 PUFA/100 g edible portion are Atlantic mackerel (2.5) and sardines in sardine oil (3.3).

Although codfish is high in omega-3 PUFA, i.e., 42.86% for the Atlantic variety and 33.33% for the Pacific variety, the levels of total fat are so low that more than 300 g/day of Atlantic codfish or more than 500 g/day of Pacific codfish would be needed to provide 1 g/day of omega-3 PUFA.

Flounder is a very popular fish in the U.S. It has 1 g of fat and 0.2 g of omega-3 PUFA per 100 g of fish. Therefore 500 g of this fish would be needed to provide 1 g of omega-PUFA. Haddock and halibut are also popular fish which are low in fat and thus require large amounts to provide 1 g/day of omega-3 PUFA. For haddock the amount of fish would be 500 g/day and for halibut the amount of fish would be 250 g/day.

For fish such as herring, 62.5 g/day of the Atlantic variety would provide 1 g/day of omega-3 PUFA; for the Pacific variety 59 g/day would suffice. Because mackerel is relatively high in fat with moderate levels of omega-3 PUFA, 40 g/day of this fish would provide 1 g/day of omega-3 PUFA and 100 g/day of the fish would provide 2.5 g/day of omega-3 PUFA. Mackerel is not a very popular fish in the U.S. Sardines packed in sardine oil provide 3.3 g omega-3 PUFA for 100 g, but the drained content of a 92 g can would provide 1.36 g omega-3 PUFA.

REGARDING SEPARATED FISH OIL IN CAPSULES

CONSUMPTION OF OMEGA-3 FATTY ACIDS

Mary G. Enig, Ph.D. 11/15/99

Omega-3 PUFA is most commonly available (usually in capsule form) from fish liver oil (cod and shark), herring oil, menhaden oil, salmon oil, and tuna oil. Capsules are usually 1 gram in size. The amounts of EPA and DHA vary depending on fish source, since these fish oils do not have identical fatty acid compositions, and also vary depending on any processing, e.g., distillation to concentrate individual fatty acids.

REGARDING REPORTS OF TOXIC LEVELS IN HUMANS

Connor & Connor (1997) recommend up to 15 g/day fish oil. Connor and colleagues fed 50 g fish oil in a 3000 kcal diet with no report of toxicity.

EXAMPLE OF RECOMMENDATIONS FROM SCIENTIFIC LITERATURE

Connor & Connor (1997) recommend dietary intake of omega-3 PUFA from fish and fish oil to prevent coronary disease in highly susceptible people and to treat patients with established coronary heart disease since the omega- PUFA "...greatly inhibit the atherosclerotic process and coronary thrombosis by many actions..."

RECOMMENDED LEVELS OF SEPARATED FISH OIL AS SUPPLEMENT

Connor & Connor (1997) recommend that fish oils be used in addition to consumption of fish at a dose range between 6 and 15 g/day if intensive treatment of various forms of hyperlipidemia as well as the production of an antithrombotic state is desired. Further for individuals who are unable to consume fish or shellfish, they advise the use of fish oil for primary prevention at levels of 2-3 g/d with the higher doses being used for secondary prevention and "the attainment of discrete endpoints of plasma lipid and lipoprotein concentrations and platelet function."

REFERENCES

Abbate R, Gori AM, Martini F, Brunelli T, Filippini M, Francalanci I, Paniccia R, Prisco D, Gensini GF, Neri Serneri GG. n-3 PUFA supplementation, monocyte PCA expression and interleukin-6 production. *Prostaglandins Leukot Essent Fatty Acids* 1996 Jun;54(6):439-44.

Abeywardena MY, Charnock JS. Dietary lipid modification of myocardial eicosanoids following ischemia and reperfusion in the rat. *Lipids* 1995 Dec;30(12):1151-6.

Albert CM, Hennekens CH, O'Donnell CJ, Ajani UA, Carey VJ, Willett WC, Ruskin JN, Manson JE. Fish consumption and risk of sudden cardiac death. *JAMA* 1998 Jan 7;279(1):23-8.

al Makdessi S, Brandle M, Ehrt M, Sweidan H, Jacob R. Myocardial protection by ischemic preconditioning: the influence of the composition of myocardial phospholipids. *Mol Cell Biochem* 1995 Apr 12;145(1):69-73.

Archer SL, Green D, Chamberlain M, Dyer AR, Liu K. Association of dietary fish and n-3 fatty acid intake with hemostatic factors in the coronary artery risk development in young adults (CARDIA) study. *Arterioscler Thromb Vasc Biol* 1998 Jul;18(7):1119-23.

Ascherio A, Rimm EB, Giovannucci EL, Spiegelman D, Stampfer M, Willett WC. Dietary fat and risk of coronary heart disease in men: cohort follow up study in the United States. *BMJ* 1996 Jul 13;313(7049):84-90.

Ascherio A, Rimm EB, Stampfer MJ, Giovannucci EL, Willett WC. Dietary intake of marine n-3 fatty acids, fish intake, and the risk of coronary disease among men. *N Engl J Med* 1995 Apr 13;332(15):977-82.

Balestrieri GP, Maffi V, Sleiman I, Spandrio S, Di Stefano O, Salvi A, Scalvini T. Fish oil supplementation in patients with heterozygous familial hypercholesterolemia. *Recenti Prog Med* 1996 Mar;87(3):102-5.

Bates EJ, Ferrante A, Smithers L, Poulos A, Robinson BS. Effect of fatty acid structure on neutrophil adhesion, degranulation and damage to endothelial cells. *Atherosclerosis* 1995 Aug;116(2):247-59.

Benistant C, Achard F, Ben Slama S, Lagarde M. Docosapentaenoic acid (22:5,n-3): metabolism and effect on prostacyclin production in endothelial cells. *Prostaglandins Leukot Essent Fatty Acids* 1996 Oct;55(4):287-92.

Billman GE, Kang JX, Leaf A. Prevention of ischemia-induced cardiac sudden death by n-3 polyunsaturated fatty acids in dogs. *Lipids* 1997 Nov;32(11):1161-8.

Mary G. Enig, Ph.D. 11/15/99

Billman GE, Kang JX, Leaf A. Prevention of sudden cardiac death by dietary pure omega-3 polyunsaturated fatty acids in dogs. *Circulation* 1999 May 11;99(18):2452-7.

Bolton-Smith C, Woodward M, Tavendale R. Evidence for age-related differences in the fatty acid composition of human adipose tissue, independent of diet. *Eur J Clin Nutr* 1997 Sep;51(9):619-24.

Bordoni A, Lopez-Jimenez JA, Spano C, Biagi P, Horrobin DF, Hrelia S. Metabolism of linoleic and alpha-linolenic acids in cultured cardiomyocytes: effect of different N-6 and N-3 fatty acid supplementation. *Mol Cell Biochem* 1996 Apr 12-26;157(1-2):217-22.

Brister SJ, Buchanan MR. Effects of linoleic acid and/or marine fish oil supplements on vessel wall thromboresistance in patients undergoing cardiac surgery. *Adv Exp Med Biol* 1997;433:275-8

Cerrato PL. Omega-3 fatty acids: nothing fishy here! *RN* 1999 Aug;62(8):59-60.

Christensen JH, Aaroe J, Knudsen N, Dideriksen K, Kornerup HJ, Dyerberg J, Schmidt EB. Heart rate variability and n-3 fatty acids in patients with chronic renal failure--a pilot study. *Clin Nephrol* 1998 Feb;49(2):102-6.

Christensen JH, Dyerberg J, Schmidt EB. n-3 fatty acids and the risk of sudden cardiac death assessed by 24-hour heart rate variability. *Lipids* 1999a;34 Suppl:S197.

Christensen JH, Christensen MS, Dyerberg J, Schmidt EB. Heart rate variability and fatty acid content of blood cell membranes: a dose-response study with n-3 fatty acids. *Am J Clin Nutr* 1999b Sep;70(3):331-7.

Christensen JH, Toft E, Christensen MS, Schmidt EB. Heart rate variability and plasma lipids in men with and without ischaemic heart disease. *Atherosclerosis* 1999c Jul;145(1):181-6

Christensen JH, Korup E, Aaroe J, Toft E, Moller J, Rasmussen K, Dyerberg J, Schmidt EB. Fish consumption, n-3 fatty acids in cell membranes, and heart rate variability in survivors of myocardial infarction with left ventricular dysfunction. *Am J Cardiol* 1997a Jun 15;79(12):1670-3.

Christensen JH, Gustenhoff P, Korup E, Aaroe J, Toft E, Moller JM, Rasmussen K, Dyerberg J, Schmidt EB. n-3 polyunsaturated fatty acids, heart rate variability and ventricular arrhythmias in post-AMI-patients. A clinical controlled trial. [Article in Danish] *Ugeskr Laeger* 1997b Sep 8;159(37):5525-9.

Mary G. Enig, Ph.D. 11/15/99

Christensen JH; Gustenhoff P; Korup E; Aaroe J; Toft E; Moller J; Rasmussen K; Dyerberg J; Schmidt EB. Effect of fish oil on heart rate variability in survivors of myocardial infarction: a double blind randomised controlled trial. *BMJ* 1996 Mar 16;312(7032):677-8.

Collie-Duguid ES, Wahle KW. Inhibitory effect of fish oil N-3 polyunsaturated fatty acids on the expression of endothelial cell adhesion molecules. *Biochem Biophys Res Commun* 1996 Mar 27;220(3):969-74.

Connor SL, Connor WE. Are fish oils beneficial in the prevention and treatment of coronary artery disease? *Am J Clin Nutr* 1997 Oct;66(4 Suppl):1020S-1031S.

Conquer JA, Holub BJ. Effect of supplementation with different doses of DHA on the levels of circulating DHA as non-esterified fatty acid in subjects of Asian Indian background. *J Lipid Res* 1998 Feb;39(2):286-92.

Conquer JA, Holub BJ. Supplementation with an algae source of docosahexaenoic acid increases (n-3) fatty acid status and alters selected risk factors for heart disease in vegetarian subjects. *J Nutr* 1996 Dec;126(12):3032-9.

Daviglus ML, Stamler J, Orenchia AJ, Dyer AR, Liu K, Greenland P, Walsh MK, Morris D, Shekelle RB. Fish consumption and the 30-year risk of fatal myocardial infarction. *N Engl J Med* 1997 Apr 10;336(15):1046-53.

De Caterina R. Fish intake and the risk of coronary disease. *N Engl J Med* 1995 Oct 5;333(14):938.

de Deckere EA, Korver O, Verschuren PM, Katan MB. Health aspects of fish and n-3 polyunsaturated fatty acids from plant and marine origin. *Eur J Clin Nutr* 1998 Oct;52(10):749-53.

Demaison L, Sergiel JP, Moreau D, Grynberg A. Influence of the phospholipid n-6/n-3 polyunsaturated fatty acid ratio on the mitochondrial oxidative metabolism before and after myocardial ischemia. *Biochim Biophys Acta* 1994 Oct 21;1227(1-2):53-9.

Enikeeva NA, Manasova PA, Kostiv ED. [Effect of diet enriched with marine omega-3 polyunsaturated fatty acids on microcirculation system in patients with cardiorespiratory pathology]. [Article in Russian] *Vopr Pitan* 1998;(4):39-41

Eritsland J, Arnesen H, Berg K, Seljeflot I, Abdelnoor M. Serum Lp(a) lipoprotein levels in patients with coronary artery disease and the influence of long-term n-3 fatty acid supplementation. *Scand J Clin Lab Invest* 1995 Jul;55(4):295-300.

Mary G. Enig, Ph.D. 11/15/99

Eritsland J, Arnesen H, Seljeflot I, Abdelnoor M, Gronseth K, Berg K, Malinow MR. Influence of serum lipoprotein(a) and homocyst(e)ine levels on graft patency after coronary artery bypass grafting. *Am J Cardiol* 1994 Dec 1;74(11):1099-102.

Eritsland J, Arnesen H, Seljeflot I, Hostmark AT. Long-term metabolic effects of n-3 polyunsaturated fatty acids in patients with coronary artery disease. *Am J Clin Nutr* 1995 Apr;61(4):831-6.

Eritsland J, Seljeflot I, Arnesen H, Westvik AB, Kierulf P. Effect of long-term, moderate-dose supplementation with omega-3 fatty acids on monocyte procoagulant activity and release of interleukin-6 in patients with coronary artery disease. *Thromb Res* 1995 Feb 15;77(4):337-46.

Ferguson JJ. Meeting highlights. Highlights of the 48th scientific sessions of the American College of Cardiology. *Circulation* 1999 Aug 10;100(6):570-5.

Freeman LM. Interventional nutrition for cardiac disease. *Clin Tech Small Anim Pract* 1998 Nov;13(4):232-7.

Fukushima T, Hojo N, Isobe A, Gao T, Shiwaku K, Yamane Y. Food intake, serum lipids and amino acids of school children in agricultural communities in Japan. *Eur J Clin Nutr* 1999 Mar;53(3):207-10.

Garg A. Treatment of diabetic dyslipidemia. *Am J Cardiol* 1998 Feb 26;81(4A):47B-51B.

Gerbi A, Barbey O, Raccach D, Coste T, Jamme I, Nouvelot A, Ouafik L, Levy S, Vague P, Maixent JM. Alteration of Na,K-ATPase isoenzymes in diabetic cardiomyopathy: effect of dietary supplementation with fish oil (n-3 fatty acids) in rats. *Diabetologia* 1997 May;40(5):496-505.

Ghafoorunissa. Dietary lipids and heart disease--the Indian context. *Natl Med J India* 1994 Nov-Dec;7(6):270-6.

[GISSI] Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. *Lancet* 1999 Aug 7;354(9177):447-55.

Goode GK, Garcia S, Heagerty AM. Dietary supplementation with marine fish oil improves in vitro small artery endothelial function in hypercholesterolemic patients: a double-blind placebo-controlled study. *Circulation* 1997 Nov 4;96(9):2802-7.

Guallar E, Aro A, Jimenez FJ, Martin-Moreno JM, Salminen I, van't Veer P, Kardinaal AF,

Gomez-Aracena J, Martin BC, Kohlmeier L, Kark JD, Mazaev VP, Ringstad J, Guillen J, Riemersma RA, Huttunen JK, Thamm M, Kok FJ. Omega-3 fatty acids in adipose tissue and risk of myocardial infarction: the EURAMIC study. *Arterioscler Thromb Vasc Biol* 1999 Apr;19(4):1111-8.

Guallar E, Hennekens CH, Sacks FM, Willett WC, Stampfer MJ. A prospective study of plasma fish oil levels and incidence of myocardial infarction in U.S. male physicians. *J Am Coll Cardiol* 1995 Feb;25(2):387-94.

Haban P, Simoncic R, Klvanova I, Ozdin L, Zidekova E. [The effect of n-3 fatty acid administration on selected indicators of cardiovascular disease risk in patients with type 2 diabetes mellitus]. [Article in Slovak] *Bratisl Lek Listy* 1998 Jan;99(1):37-42.

Hamazaki T, Sawazaki S, Itomura M, Asaoka E, Nagao Y, Nishimura N, Yazawa K, Kuwamori T, Kobayashi M. The effect of docosahexaenoic acid on aggression in young adults. A placebo-controlled double-blind study. *J Clin Invest* 1996 Feb 15;97(4):1129-33.

Hardy SC, Kleinman RE. Fat and cholesterol in the diet of infants and young children: implications for growth, development, and long-term health. *J Pediatr* 1994 Nov;125(5 Pt 2):S69-77.

Harris WS. n-3 fatty acids and human lipoprotein metabolism: an update. *Lipids* 1999;34 Suppl:S257-8.

Harris WS. Nonpharmacologic treatment of hypertriglyceridemia: focus on fish oils. *Clin Cardiol* 1999 Jun;22(6 Suppl):II40-3.

Harris WS. Dietary fish oil and blood lipids. *Curr Opin Lipidol* 1996 Feb;7(1):3-7.

Harris WS, Ginsberg HN, Arunakul N, Shachter NS, Windsor SL, Adams M, Berglund L, Osmundsen K. Safety and efficacy of Omacor in severe hypertriglyceridemia. *J Cardiovasc Risk* 1997 Oct-Dec;4(5-6):385-91.

Hartman IS. alpha-Linolenic acid: a preventive in secondary coronary events? *Nutr Rev* 1995 Jul;53(7):194-7.

Hayashi M, Nasa Y, Tanonaka K, Sasaki H, Miyake R, Hayashi J, Takeo S. The effects of long-term treatment with eicosapentaenoic acid and docosahexaenoic acid on hypoxia/reoxygenation injury of isolated cardiac cells in adult rats. *J Mol Cell Cardiol* 1995 Sep;27(9):2031-41.

Heemskerk JW, Vossen RC, van Dam-Mieras MC. Polyunsaturated fatty acids and function of platelets and endothelial cells. *Curr Opin Lipidol* 1996 Feb;7(1):24-9.

Mary G. Enig, Ph.D. 11/15/99

Henzen C. [Fish oil--healing principle in the Eskimo diet]? [Article in German] *Schweiz Rundsch Med Prax* 1995 Jan 3;84(1):11-5.

Herrmann W, Biermann J, Kostner GM. Comparison of effects of N-3 to N-6 fatty acids on serum level of lipoprotein(a) in patients with coronary artery disease. *Am J Cardiol* 1995 Sep 1;76(7):459-62.

Hishinuma T, Yamazaki T, Mizugaki M. Effects of long-term supplementation of eicosapentanoic and docosahexanoic acid on the 2-, 3-series prostacyclin production by endothelial cells. *Prostaglandins Other Lipid Mediat* 1999 Jul;57(5-6):333-40.

Hojo N, Fukushima T, Isobe A, Gao T, Shiwaku K, Ishida K, Ohta N, Yamane Y. Effect of serum fatty acid composition on coronary atherosclerosis in Japan. *Int J Cardiol* 1998 Sep 1;66(1):31-8.

Horrocks LA, Yeo YK. Health benefits of docosahexaenoic acid. *Pharmacol Res* 1999 Sep;40(3):211-25.

Hrboticky N, Sellmayer A. [Nutrition in prevention of coronary heart disease]. [Article in German] *Z Arztl Fortbild (Jena)* 1996 Feb;90(1):11-8.

Hu FB, Stampfer MJ, Manson JE, Rimm EB, Wolk A, Colditz GA, Hennekens CH, Willett WC. Dietary intake of alpha-linolenic acid and risk of fatal ischemic heart disease among women. *Am J Clin Nutr* 1999 May;69(5):890-7.

Hwang DH, Chanmugam PS, Ryan DH, Boudreau MD, Windhauser MM, Tulley RT, Brooks ER, Bray GA. Does vegetable oil attenuate the beneficial effects of fish oil in reducing risk factors for cardiovascular disease? *Am J Clin Nutr* 1997 Jul;66(1):89-96.

Isensee H, Jacob R. Differential effects of various oil diets on the risk of cardiac arrhythmias in rats. *J Cardiovasc Risk* 1994 Dec;1(4):353-9.

Johansen O, Brekke M, Seljeflot I, Abdelnoor M, Arnesen H. N-3 fatty acids do not prevent restenosis after coronary angioplasty: results from the CART study. Coronary Angioplasty Restenosis Trial. *J Am Coll Cardiol* 1999 May;33(6):1619-26.

Johansen O, Seljeflot I, Hostmark AT, Arnesen H. The effect of supplementation with omega-3 fatty acids on soluble markers of endothelial function in patients with coronary heart disease. *Arterioscler Thromb Vasc Biol* 1999 Jul;19(7):1681-6.

Mary G. Enig, Ph.D. 11/15/99

Kaminskas A, Zieden B, Elving B, Kristenson M, Abaravicius A, Bergdahl B, Olsson AG, Kucinskiene Z. Adipose tissue fatty acids in men from two populations with different cardiovascular risk: the LiVicordia study. *Scand J Clin Lab Invest* 1999 May;59(3):227-32.

Kang JX, Leaf A. The cardiac antiarrhythmic effects of polyunsaturated fatty acid. *Lipids* 1996 Mar;31 Suppl: S41-4.

Kang JX, Leaf A. Prevention and termination of beta-adrenergic agonist-induced arrhythmias by free polyunsaturated fatty acids in neonatal rat cardiac myocytes. *Biochem Biophys Res Commun* 1995 Mar 17;208(2):629-36.

Katan MB. Fish and heart disease. *N Engl J Med* 1995 Apr 13;332(15):1024-5.

Katan MB. Fish and heart disease: what is the real story? *Nutr Rev* 1995 Aug;53(8):228-30.

Kinoshita I, Itoh K, Nishida-Nakai M, Hirota H, Otsuji S, Shibata N. Antiarrhythmic effects of eicosapentaenoic acid during myocardial infarction--enhanced cardiac microsomal (Ca(2+)-Mg2+)-ATPase activity. *Jpn Circ J* 1994 Dec;58(12):903-12.

Kothny W, Angerer P, Stork S, von Schacky C. Short term effects of omega-3 fatty acids on the radial artery of patients with coronary artery disease. *Atherosclerosis* 1998 Sep;140(1):181-6.

Kozycheva EV, Slezka IE. [Lipid-correcting effect of marine unsaturated fatty acids for prophylaxis of cardiovascular diseases in boys]. [Article in Russian] *Vopr Pitan* 1998;(4):27-31.

Kromhout D, Feskens EJ, Bowles CH. The protective effect of a small amount of fish on coronary heart disease mortality in an elderly population. *Int J Epidemiol* 1995 Apr;24(2):340-5.

Ku K, Nosaka S, Hashimoto M, Kin S, Saitoh Y, Alam MS, Masumura S, Nakayama K. The effects of omega-3 polyunsaturated (correction of polyunsaturated) fatty acids on the recovery of cardiac function following cold preservation and reperfusion in hyperlipidemic rats. *Transplantation* 1996 Sep 27;62(6):735-42.

Landmark K. [Fish, fish oils, arrhythmias and sudden death]. [Article in Norwegian] *Tidsskr Nor Laegeforen* 1998 Jun 10;118(15):2328-31.

Lavie CJ, Milani RV, Ventura HO, Messerli FH, Murgo JP. Cardiac rehabilitation, exercise training, and preventive cardiology research at Ochsner Heart and Vascular Institute. *Tex Heart Inst J* 1995;22(1):44-52.

Leaf A. Dietary prevention of coronary heart disease: the Lyon Diet Heart Study. *Circulation* 1999 Feb 16;99(6):733-5.

Mary G. Enig, Ph.D. 11/15/99

Leaf A. Fish intake and the risk of coronary disease. *N Engl J Med* 1995 Oct 5;333(14):937-8.

Leaf A. Omega-3 fatty acids and prevention of ventricular fibrillation. *Prostaglandins Leukot Essent Fatty Acids* 1995 Feb-Mar;52(2-3):197-8.

Leaf A, Kang JX. Dietary n-3 fatty acids in the prevention of lethal cardiac arrhythmias. *Curr Opin Lipidol* 1997 Feb;8(1):4-6.

Leaf A, Kang JX. Prevention of cardiac sudden death by N-3 fatty acids: a review of the evidence. *J Intern Med* 1996 Jul;240(1):5-12.

Leaf A, Kang JX, Xiao YF, Billman GE. n-3 fatty acids in the prevention of cardiac arrhythmias. *Lipids* 1999;34 Suppl:S187-9.

Li D, Sinclair A, Wilson A, Nakkote S, Kelly F, Abedin L, Mann N, Turner A. Effect of dietary alpha-linolenic acid on thrombotic risk factors in vegetarian men. *Am J Clin Nutr* 1999 May;69(5):872-82.

Li Y, Kang JX, Leaf A. Differential effects of various eicosanoids on the production or prevention of arrhythmias in cultured neonatal rat cardiac myocytes. *Prostaglandins* 1997 Aug;54(2):511-30.

Liutova LV, Karabasova MA, Andreenko GV, Panchenko VM, Isaev VA, Sorokaletov SM, Ershov AA. [Effect of docanol on the status of hemostasis and fibrinolysis systems, as well as lipid spectrum in patients with ischemic heart disease depending on the initial level of fibrinolytic activity]. [Article in Russian] *Vopr Med Khim* 1996 Jan-Mar;42(1):64-70.

Mantzioris E, James MJ, Gibson RA, Cleland LG. Differences exist in the relationships between dietary linoleic and alpha-linolenic acids and their respective long-chain metabolites. *Am J Clin Nutr* 1995 Feb;61(2):320-4.

Marckmann P, Gronbaek M. Fish consumption and coronary heart disease mortality. A systematic review of prospective cohort studies. *Eur J Clin Nutr* 1999 Aug;53(8):585-90.

Masley SC. Dietary therapy for preventing and treating coronary artery disease. *Am Fam Physician* 1998 Mar 15;57(6):1299-1306, 1307-9.

McLennan PL, Raederstorff D. Diabetes puts myocardial n-3 fatty acid status at risk in the absence of supplementation in the rat. *Lipids* 1999;34 Suppl:S91-2.

Mary G. Enig, Ph.D. 11/15/99

McLennan P, Howe P, Abeywardena M, Muggli R, Raederstorff D, Mano M, Rayner T, Head R. The cardiovascular protective role of docosahexaenoic acid. *Eur J Pharmacol* 1996 Apr 4;300(1-2):83-9.

Mensink R. Essential and trans-fatty acids. *Curr Opin Lipidol* 1996 Feb;7(1):1-2.

Miller GJ. Effects of diet composition on coagulation pathways. *Am J Clin Nutr* 1998 Mar;67(3 Suppl):542S-545S.

Mizutani M, Asano M, Roy S, Nakajima T, Soma M, Yamashita K, Okuda Y. Omega-3 polyunsaturated fatty acids inhibit migration of human vascular smooth muscle cells in vitro. *Life Sci* 1997;61(19):PL269-74.

Moller JM, Nielsen GL, Ekelund S, Schmidt EB, Dyerberg J. Homocysteine in Greenland Inuits. *Thromb Res* 1997 May 15;86(4):333-5.

Mori TA, Beilin LJ, Burke V, Morris J, Ritchie J. Interactions between dietary fat, fish, and fish oils and their effects on platelet function in men at risk of cardiovascular disease. *Arterioscler Thromb Vasc Biol* 1997 Feb;17(2):279-86.

Morris MC, Manson JE, Rosner B, Buring JE, Willett WC, Hennekens CH. Fish consumption and cardiovascular disease in the physicians' health study: a prospective study. *Am J Epidemiol* 1995 Jul 15;142(2):166-75.

Mutanen M, Freese R. Polyunsaturated fatty acids and platelet aggregation. *Curr Opin Lipidol* 1996 Feb;7(1):14-9.

Nair SS, Leitch J, Falconer J, Garg ML. Cardiac (n-3) non-esterified fatty acids are selectively increased in fish oil-fed pigs following myocardial ischemia. *J Nutr* 1999 Aug;129(8):1518-23.

Nair SS, Leitch JW, Falconer J, Garg ML. Prevention of cardiac arrhythmia by dietary (n-3) polyunsaturated fatty acids and their mechanism of action. *J Nutr* 1997 Mar;127(3):383-93.

Nakamura N, Hamazaki T, Ohta M, Okuda K, Urakaze M, Sawazaki S, Yamazaki K, Satoh A, Temaru R, Ishikura Y, Takata M, Kishida M, Kobayashi M. Joint effects of HMG-CoA reductase inhibitors and eicosapentaenoic acids on serum lipid profile and plasma fatty acid concentrations in patients with hyperlipidemia. *Int J Clin Lab Res* 1999;29(1):22-5.

Nasa Y, Hayashi M, Sasaki H, Hayashi J, Takeo S. Long-term supplementation with eicosapentaenoic acid salvages cardiomyocytes from hypoxia/reoxygenation-induced injury in rats fed with fish-oil-deprived diet. *Jpn J Pharmacol* 1998 Jun;77(2):137-46.

Nordoy A. Dietary fatty acids and coronary heart disease. *Lipids* 1999;34 Suppl:S19-22.

Nordoy A, Bonna KH, Nilsen H, Berge RK, Hansen JB, Ingebretsen OC. Effects of Simvastatin and omega-3 fatty acids on plasma lipoproteins and lipid peroxidation in patients with combined hyperlipidaemia. *J Intern Med* 1998 Feb;243(2):163-70.

Novgorodtseva TP, Endakova EA, Kozycheva EV, Kitaiskaia LS. [Modification to the erythrocyte membrane lipids in patients with cardiovascular pathology during balneotherapy]. [Article in Russian] *Vopr Kurortol Fizioter Lech Fiz Kult* 1999 Jan-Feb;(1):12-5.

O'Farrell S, Jackson MJ. Dietary polyunsaturated fatty acids, vitamin E and hypoxia/reoxygenation-induced damage to cardiac tissue. *Clin Chim Acta* 1997 Nov 28;267(2):197-211.

Opstvedt J. Fish lipids: more than n-3 fatty acids? *Med Hypotheses* 1997 Jun;48(6):481-3.

Pakala R, Pakala R, Sheng WL, Benedict CR. Serotonin fails to induce proliferation of endothelial cells preloaded with eicosapentaenoic acid and docosahexaenoic acid. *Atherosclerosis* 1999 Jul;145(1):137-46.

Panchenko VM, Nazlukhanian SO, Isaev VA, Ershov AA, Chernova GI, Liutova LV, Andreenko GV, Dzhnashvili PK. [Effects of eifitol on lipid metabolism in patients with clinically different ischemic heart diseases]. [Article in Russian] *Klin Med (Mosk)* 1997;75(3):39-41.

Pepe S, McLennan PL. Dietary fish oil confers direct antiarrhythmic properties on the myocardium of rats. *J Nutr* 1996 Jan;126(1):34-42.

Pitt B. Results of recent trials on the progression of coronary artery disease and recurrent ischemic events: implications for interventional cardiology. *J Interv Cardiol* 1994 Dec;7(6):515-8.

Pogozheva AV, Rozanova IA, Miagkova MA, Sorokovoi KV, Panchenko ON, Trubacheva ZhN [Comparative study on the effect of antiatherosclerotic diet enriched with polyunsaturated omega-3 fatty acids of plant and animal origin on the level of natural antibodies against catecholamines in patients with cardiovascular diseases]. [Article in Russian] *Vopr Pitan* 1998;(4):32-5.

Pogozheva AV, Rozanova IA, Sorokovoi KV, Karagodina ZV, Levachev MM. [Lipid peroxidation in patients with ischemic heart disease, hyperlipidemia and/or hypertension while polyunsaturated omega-3 fatty acids of plant and animal origin were in their diet]. [Article in Russian] *Vopr Pitan* 1997;(4):32-5.

Mary G. Enig, Ph.D. 11/15/99

Pogozheva AV, Samsonov MA, Levachev MM. [Clinico-biochemical validation of the use of PUFA omega-3 in patients with ischemic heart disease, familial hyperlipoproteinemias and hypertension]. [Article in Russian] *Vopr Pitan* 1996;(1):34-6.

Ponte E, Cafagna D, Balbi M. [Cardiovascular disease and omega-3 fatty acids]. [Article in Italian] *Minerva Med* 1997 Sep;88(9):343-53.

Reithmann C, Scheininger C, Bulgan T, Werdan K. Exposure to the n-3 polyunsaturated fatty acid docosahexaenoic acid impairs alpha 1-adrenoceptor-mediated contractile responses and inositol phosphate formation in rat cardiomyocytes. *Naunyn Schmiedebergs Arch Pharmacol* 1996 Jul;354(2):109-19.

Roche HM. Unsaturated fatty acids. *Proc Nutr Soc* 1999 May;58(2):397-401.

Roche HM, Gibney MJ. Long-chain n-3 polyunsaturated fatty acids and triacylglycerol metabolism in the postprandial state. *Lipids* 1999;34 Suppl:S259-65.

Rodrigo GC, Dhanapala S, Macknight AD. Effects of eicosapentaenoic acid on the contraction of intact, and spontaneous contraction of chemically permeabilized mammalian ventricular myocytes. *J Mol Cell Cardiol* 1999 Apr;31(4):733-43.

Rozanova IA, Pogozheva AV, Miagkova MA, Savitskaia IuA. [Effect of anti-atherosclerotic diet with polyunsaturated fatty acids omega-3 from linseed oil on dynamics of natural antibodies to bradykinin and angiotensin II in blood serum of patients with cardiovascular diseases]. [Article in Russian] *Vopr Pitan* 1998;(2):32-4

Rozanova IA, Pogozheva AV, Kupakova SN, Lupinovich VL, Karagodina ZV, Levachev MM, Samsonov MA. [Effect of antiatherosclerotic diet, containing polyunsaturated fatty acids of the omega-3 family from flax oil, on fatty acid composition of cell membranes of patients with ischemic heart disease. Hypertensive disease and hyperlipoproteinemia]. [Article in Russian] *Vopr Pitan* 1997;(5):15-7.

Sacks FM, Stone PH, Gibson CM, Silverman DI, Rosner B, Pasternak RC. Controlled trial of fish oil for regression of human coronary atherosclerosis. HARP Research Group. *J Am Coll Cardiol* 1995 Jun;25(7):1492-8.

Salonen JT, Nyyssonen K, Salonen R. Fish intake and the risk of coronary disease. *N Engl J Med* 1995 Oct 5;333(14):937; discussion 938.

Samsonov MA, Pogozheva AV. [Marine fish oil in the prevention and treatment of cardiovascular diseases]. [Article in Russian] *Vestn Ross Akad Med Nauk* 1996;(12):43-9.

Mary G. Enig, Ph.D. 11/15/99

Sanchez-Muniz FJ, Bastida S, Viejo JM, Terpstra AH. Small supplements of N-3 fatty acids change serum low density lipoprotein composition by decreasing phospholipid and apolipoprotein B concentrations in young adult women. *Eur J Nutr* 1999 Feb;38(1):20-7.

Sanders TA. Effects of unsaturated fatty acids on blood clotting and fibrinolysis. *Curr Opin Lipidol* 1996 Feb;7(1):20-3.

Santos MJ, Llopis J, Mataix FJ, Urbano G, Lopez Jurado M. Influence of dietary fish on fatty acid composition of the erythrocyte membrane in coronary heart disease patients. *Int J Vitam Nutr Res* 1996;66(4):378-85.

Santos MJ, Lopez-Jurado M, Llopis J, Urbano G, Mataix FJ. Influence of dietary supplementation with fish on plasma fatty acid composition in coronary heart disease patients. *Ann Nutr Metab* 1995;39(1):52-62.

Schaefer EJ, Lichtenstein AH, Lamon-Fava S, Contois JH, Li Z, Goldin BR, Rasmussen H, McNamara JR, Ordovas JM. Effects of National Cholesterol Education Program Step 2 diets relatively high or relatively low in fish-derived fatty acids on plasma lipoproteins in middle-aged and elderly subjects. *Am J Clin Nutr* 1996 Feb;63(2):234-41.

Schindler OS, Rost R. [Effect of low dose omega-3 fatty acid supplementation on plasma lipids and lipoproteins in coronary patients with dyslipoproteinemia]. [Article in German] *Z Ernahrungswiss* 1996 Jun;35(2):191-8.

Schmidt EB. n-3 fatty acids and the risk of coronary heart disease. *Dan Med Bull* 1997 Feb;44(1):1-22.

Schmidt EB, Dyerberg J. n-3 fatty acids and coronary heart disease--the urgent need of clinical trials. *Lipids* 1999;34 Suppl:S303-5.

Schmidt EB, Varming K, Moller JM, Bulow Pedersen I, Madsen P, Dyerberg J. No effect of a very low dose of n-3 fatty acids on monocyte function in healthy humans. *Scand J Clin Lab Invest* 1996 Feb;56(1):87-92.

Seidelin KN. Fatty acid composition of adipose tissue in humans. Implications for the dietary fat-serum cholesterol-CHD issue. *Prog Lipid Res* 1995;34(3):199-217.

Sergiel JP, Martine L, Raederstorff D, Grynberg A, Demaison L. Individual effects of dietary EPA and DHA on the functioning of the isolated working rat heart. *Can J Physiol Pharmacol* 1998 Jul-Aug;76(7-8):728-36

Mary G. Enig, Ph.D. 11/15/99

Severus WE, Ahrens B, Stoll AL. Omega-3 fatty acids--the missing link? *Arch Gen Psychiatry* 1999 Apr;56(4):380-1.

Sheard NF. Fish consumption and risk of sudden cardiac death. *Nutr Rev* 1998 Jun;56(6):177-9.

Shimizu T. The future potential of eicosanoids and their inhibitors in paediatric practice. *Drugs* 1998 Aug;56(2):169-76.

Skuladottir GV, Gudmundsdottir S, Olafsson GB, Sigurdsson SB, Sigfusson N, Axelsson J. Plasma fatty acids and lipids in two separate, but genetically comparable, Icelandic populations. *Lipids* 1995 Jul;30(7):649-55.

Siguel EN. Dietary sources of long-chain n-3 polyunsaturated fatty acids. *JAMA* 1996 Mar 20;275(11):836-7.

Simon JA, Hodgkins ML, Browner WS, Neuhaus JM, Bernert JT Jr, Hulley SB. Serum fatty acids and the risk of coronary heart disease. *Am J Epidemiol* 1995 Sep 1;142(5):469-76.

Singh RB, Niaz MA, Sharma JP, Kumar R, Rastogi V, Moshiri M. Randomized, double-blind, placebo-controlled trial of fish oil and mustard oil in patients with suspected acute myocardial infarction: the Indian experiment of infarct survival--4. *Cardiovasc Drugs Ther* 1997 Jul;11(3):485-91.

Sirtori CR, Paoletti R, Mancini M, Crepaldi G, Manzato E, Rivellesse A, Pamparana F, Stragliotto E. N-3 fatty acids do not lead to an increased diabetic risk in patients with hyperlipidemia and abnormal glucose tolerance. Italian Fish Oil Multicenter Study. *Am J Clin Nutr* 1997 Jun;65(6):1874-81.

Siscovick DS, Raghunathan TE, King I, Weinmann S, Wicklund KG, Albright J, Bovbjerg V, Arbogast P, Smith H, Kushi LH, et al. Dietary intake and cell membrane levels of long-chain n-3 polyunsaturated fatty acids and the risk of primary cardiac arrest. *JAMA* 1995 Nov 1;274(17):1363-7.

Sorokovoi KV, Pogozheva AV, Kulakova SN, Lupinovich VL, Karagodina ZV, Levachev MM, Samsonov MA, Vorsanovich EA. [Study of metabolic and clinical effects of polyunsaturated fatty acids omega-3 from "Eicovit" in patients with ischemic heart disease and familial hyperlipoproteinemia]. [Article in Russian] *Vopr Pitan* 1998;(2):29-31.

Sorokovoi KV, Pogozheva AV, Sergeeva KV. [Effect of diet including omega-3 polyunsaturated fatty acids on the status of T-cell immunity in patients with ischemic heart disease and hyperproteinemia]. [Article in Russian] *Vopr Pitan* 1997;(5):6-8.

Starkopf J, Andreassen TV, Bugge E, Ytrehus K. Lipid peroxidation, arachidonic acid and products of the lipoxygenase pathway in ischaemic preconditioning of rat heart. *Cardiovasc Res* 1998 Jan;37(1):66-75.

Stone NJ. Fish consumption, fish oil, lipids, and coronary heart disease. *Am J Clin Nutr* 1997 Apr;65(4):1083-6.

Stone NJ. Fish consumption, fish oil, lipids, and coronary heart disease. *Circulation* 1996 Nov 1;94(9):2337-40.

Supari F, Ungerer T, Harrison DG, Williams JK. Fish oil treatment decreases superoxide anions in the myocardium and coronary arteries of atherosclerotic monkeys. *Circulation* 1995 Feb 15;91(4):1123-8.

Suzuki H, Manabe S, Wada O, Crawford MA. Rapid incorporation of docosahexaenoic acid from dietary sources into brain microsomal, synaptosomal and mitochondrial membranes in adult mice. *Int J Vitam Nutr Res* 1997;67(4):272-8.

Tagawa H, Shimokawa H, Tagawa T, Kuroiwa-Matsumoto M, Hirooka Y, Takeshita A. Long-term treatment with eicosapentaenoic acid augments both nitric oxide-mediated and non-nitric oxide-mediated endothelium-dependent forearm vasodilatation in patients with coronary artery disease. *J Cardiovasc Pharmacol* 1999 Apr;33(4):633-40.

Takeo S, Nasa Y, Tanonaka K, Yabe K, Nojiri M, Hayashi M, Sasaki H, Ida K, Yanai K. Effects of long-term treatment with eicosapentaenoic acid on the heart subjected to ischemia/reperfusion and hypoxia/reoxygenation in rats. *Mol Cell Biochem* 1998 Nov;188(1-2):199-208.

Temple NJ. Dietary fats and coronary heart disease. *Biomed Pharmacother* 1996;50(6-7):261-8.

Thompson RL, Pyke SD, Scott EA, Thompson SG, Wood DA. Dietary change after smoking cessation: a prospective study. *Br J Nutr* 1995 Jul;74(1):27-38.

Tinker LF, Parks EJ, Behr SR, Schneeman BO, Davis PA. (n-3) fatty acid supplementation in moderately hypertriglyceridemic adults changes postprandial lipid and apolipoprotein B responses to a standardized test meal. *J Nutr* 1999 Jun;129(6):1126-34.

Tsuruta K, Ogawa H, Yasue H, Sakamoto T, Miyao Y, Tanae H, Kaiga K. Effect of purified eicosapentaenoate ethyl ester on fibrinolytic capacity in patients with stable coronary artery disease and lower extremity ischaemia. *Coron Artery Dis* 1996 Nov;7(11):837-42.

Mary G. Enig, Ph.D. 11/15/99

Turley E, Wallace JM, Gilmore WS, Strain JJ. Fish oil supplementation with and without added vitamin E differentially modulates plasma antioxidant concentrations in healthy women. *Lipids* 1998 Dec;33(12):1163-7.

von Schacky C, Angerer P, Kothny W, Theisen K, Mudra H. The effect of dietary omega-3 fatty acids on coronary atherosclerosis. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 1999 Apr 6;130(7):554-62.

Voskuil DW, Feskens EJM, Katan MB, Kromhout D. Intake and sources of alpha-linolenic acid in Dutch elderly men. *Eur J Clin Nutr* 1996 Dec;50(12):784-7.

Yamamoto H, Yoshimura H, Noma M, Suzuki S, Kai H, Tajimi T, Sugihara M, Kikuchi Y. Improvement of coronary vasomotion with eicosapentaenoic acid does not inhibit acetylcholine-induced coronary vasospasm in patients with variant angina. *Jpn Circ J* 1995 Sep;59(9):608-16. Published erratum appears in *Jpn Circ J* 1995 Nov;59(11):773-4.

Weintraub M, Charach G, Grosskopf I. Disturbances in dietary fat metabolism and their role in the development of atherosclerosis. *Biomed Pharmacother* 1997;51(8):311-3.

Zak A, Zeman M, Tvrzicka E. [Effect of fish oils on plasma lipid risk factors and esterified fatty acids in primary hyperlipoproteinemia]. [Article in Czech] *Sb Lek* 1997;98(3):213-24.

Zak A, Zeman M, Vitkova D. [Comparison of the effects of omega-3 polyunsaturated fatty acids and Olbetam (Acipimox) in the treatment of hypertriglyceridemia]. [Article in Czech] *Cas Lek Cesk* 1994 Dec 22;133(24):755-8.

Zampelas A, Peel AS, Gould BJ, Wright J, Williams CM. Polyunsaturated fatty acids of the n-6 and n-3 series: effects on postprandial lipid and apolipoprotein levels in healthy men. *Eur J Clin Nutr* 1994 Dec;48(12):842-8.

[No author] Fish and the heart. *Harv Health Lett* 1998 Mar;23(5):8.

[No author] Fish and heart health. Make fish a regular on your plate. *Mayo Clin Health Lett* 1998 Jul;16(7):7.